

YUMS - 98/05
February 1997

IMPLICATIONS OF THE HERA EVENTS FOR THE R-PARITY BREAKING SUSY SIGNALS AT TEVTARON

Monoranjana Guchait

Department of Physics, Yonsei University, Seoul 120-749, South Korea

I report here on the systematic analysis at Tevatron of the R-parity violating SUSY signals which correspond to the possible solution of anomalous HERA events

The two experimental groups at DESY, H1 and ZEUS, have reported some anomalous high Q^2 events from the e^+p collider. The H1 has seen 12 neutral current events, at $Q^2 > 15,000 \text{ GeV}^2$ against the Standard Model(SM) prediction of 5 events¹; while the ZEUS has reported 5 events at $Q^2 > 20,000 \text{ GeV}^2$ against the prediction² of 2. The excess of 7 events observed by H1 has clustered around a common e^+q mass of

$$M \simeq 200 \text{ GeV} \quad (1)$$

which is inconsistent to ZEUS events³. These are based on 1994-'96 data and corresponding to a combined luminosity 34 pb^{-1} for the two experiments. Now taking together, there are 10 excess events which corresponds to a cross section

$$\sigma \simeq 0.4 \text{ pb} \quad (2)$$

for the 80% detection efficiency for each experiments.

Many mechanisms of new physics was proposed to explain these^{4,5,?,7} new events. Among the proposed mechanisms of new physics, the R-parity violating(RPV) SUSY model is the most popular one. The purpose of this work⁸ is to study the implications of the RPV SUSY model at Tevatron

The Minimal Supersymmetric Standard Model(MSSM) with explicit R-parity breaking⁹ contains the following Yukawa terms in the Lagrangian

$$L = \lambda_{ijk} l_i \tilde{l}_j \bar{e}_k + \lambda'_{ijk} l_i \tilde{q}_j \bar{d}_k + \lambda''_{ijk} \bar{d}_i \tilde{d}_j \bar{u}_k, \quad (3)$$

plus analogous terms from the permutation of the tilde, denoting the scalar superpartner. We have used the same notation as of ref⁸. The terms relevant for the HERA events are

$$\lambda'_{1jk} (l_1 \tilde{q}_j \bar{d}_k + l_1 q_j \tilde{d}_k) + h.c. \quad (4)$$

Therefore, for a given quark flux one can easily calculate the cross section for these process at HERA when $\lambda'\sqrt{B}$ is given, where B denotes the squark branching fraction into the shown channel. While the size $\lambda'\sqrt{B}$ is constrained by HERA data as given by Eq. 2 and the value of λ' is restricted by its upper bounds from some other process^{5,8}. It is shown^{4,5} that the viable couplings which are allowed by the HERA data and as well as by its corresponding existing upper limits are the λ'_{121} and λ'_{131} which stand for the scharm and stop squark production from a valence quark i.e .

$$e^+d \rightarrow \tilde{c}_L \rightarrow e^+d. \quad (5)$$

$$e^+d \rightarrow \tilde{t}_L \rightarrow e^+d, \quad (6)$$

where the L subscript stands for left chirality. Therefore, we have concentrated only on these two scenarios and called them scharm and stop scenario for corresponding to the λ'_{121} and λ'_{131} couplings respectively.

Since the product of $\lambda'\sqrt{B}$ is constrained by the cross section, Eq. 2, of the HERA events, it is possible to express B in terms of the MSSM parameters under the assumption that there is only one dominant RPV coupling⁸. The dependence of B on the MSSM parameter space for a given scenario will be discussed below.

It is to be noted that in our analysis we assume a common gaugino and a common sfermion mass at the unification scale which results the masses of gauginos ($\tilde{g}, \tilde{W}$ and $\tilde{B}$) are related via their gauge couplings at the electroweak unification scale⁸. Under this assumption we have investigated the possibility of finding the stop and scharm scenario at the Tevatron collider. The results are in order.

0.1 The Stop Scenario at Tevatron:

The dominant production mechanism for stop at Tevatron are the leading order QCD process of quark-antiquark and gluon-gluon fusion¹⁰

$$\bar{q}q \rightarrow \tilde{t}_1\tilde{t}_1 \quad , \quad gg \rightarrow \tilde{t}_1\tilde{t}_1. \quad (7)$$

Because of the negligible contribution of the RPV couplings, we have not considered it here.

The dominant decay mode of stop is the RPV mode

$$\tilde{t}_1 \rightarrow e^+d. \quad (8)$$

as long as $M_{\tilde{t}_1} \simeq M_{\tilde{W}_1}$, because the decay mode in the neutralino channel is higher order process due to the large top quark mass. But the R-parity

conserving decay mode

$$\tilde{t}_1 \rightarrow b\tilde{W}_i \quad (9)$$

will dominate when it will be kinematically allowed. In Fig.1, the values of B




Figure 1: The stop branching fraction (B) is shown as a contour plot in the M_2, μ plane for the stop mass $M_{\tilde{t}_1} = 200\text{GeV}$, $\tan\beta = 2$ and $\theta_{\tilde{t}} = 0^\circ$. Contours for the lighter chargino mass of 180GeV(solid line) and 85GeV(dotted line) are shown.

are shown over the relevant MSSM parameter space. It is clear from the fig.1 that the the signal of stop at Tevatron depends very crucially on the MSSM parameter space. In our analysis, we have considered three types of decay configuration of stop which are discussed below.

(a) Direct Leptonic decay: In this case both the stop squarks decay in the RPV channel(Eq. 8) resulting a pair of hard and isolated dilepton along with a pair of jets. The corresponding 95% C.L. limits of 3 events as obtained from the 1st generation of Leptoquark search gives a mass bound by CDF $M > 210\text{ GeV}$ ¹¹ where as this value for the D0 group¹² is $M > 225\text{GeV}$. Thus CDF and D0 leptoquark limits would rule out the stop scenario if the direct leptonic decays are dominant, i.e. $M_{\tilde{W}_1} \geq M_{\tilde{t}_1}$. However, dilepton cross section for the $B \leq 0.2$ is so small that it is not possible to probe them in the present data

set of CDF and D0. But in the MI run one can probe the signal cross section down to $B = 0.2$. Nonetheless, there is a significant MSSM parameter space where $B \leq 0.2$ and it is possible to probe this parameter space in the MI run. But it may be possible to probe this entire MSSM parameter space at TEV33 where luminosity will be 20 fb^{-1} .

(b)Mixed Mode: This mode corresponds to direct decay of one stop (eq. 8) and other undergoes cascade decay as

$$\tilde{t}_1 \rightarrow b\tilde{W}_1, \tilde{W}_1 \rightarrow q'\bar{q}\tilde{Z}_1, \tilde{Z}_1 \xrightarrow{\lambda'_{131}} \bar{b}\bar{\nu}d + h.c. \quad (10)$$

Thus the final states consists of a very hard $e^\pm$, a large number of jets including a pair of b and a modest amount of missing p_T carried by the neutrino. The dominant background to this process is from $t\bar{t}$. Fig.2 shows the signal cross section for a stop mass 200 GeV along with the $t\bar{t}$ background at $\sqrt{s} = 2\text{TeV}$ for a following MSSM parameters,

$$M_2 = 150\text{GeV}, \mu = -400\text{GeV}, \tan\beta = 2 \Rightarrow M_{\tilde{W}_1} = 158\text{GeV}, M_{\tilde{Z}_1} = 77\text{GeV}; \quad (11)$$

Note that this signal rate is quite insensitive of MSSM parameters. From Fig.2 it is clear that the signal can be separated from the background by exploiting the clustering of invariant mass of the electron with the hardest jets at $M \simeq 200\text{GeV}$. The size of the signal in Fig.2 is about 50fb which means few events are expected for the MI run for the MSSM parameter space (in Fig.1) where the branching fraction is $B \simeq 7\%$.

(c)Cascade decays: It corresponds to the cascade decays, eq. 10 ,of both the stop squarks resulting in the final states four b quarks and a missing p_T carried by the neutrinos. We need to do the tripple b tagging to suppress background which is mainly from $t\bar{t}$ and $b\bar{b}$ production. We have estimated the signal cross section for two different channels (i) 3 b quarks + missing p_T and (ii) 1 lepton + 3 b + missing p_T . We found no viable signature for detecting the stop pair production for the integrated luminosity 2 fb^{-1} . But in the MI run it may be possible to probe entire MSSM parameter space for larger b-tagging efficiency.

0.2 The Charm Squark(Scharm!) Scenario at Tevatron

In this case one expects 8 species of roughly degenerate squarks along with a gluino of comaparable or smaller mass. Only one of them, the left handed charm squark $\tilde{c}_L$, has the R violating decay mode

$$\tilde{c}_L \rightarrow e^+d. \quad (12)$$




Figure 2: The signal cross section(solid line) corresponding to the mixed mode(b) is shown for stop mass of 200 GeV along with the $t\bar{t}$ background(dashed line) against the p_T of the electron at 2 TeV. The MSSM parameters are $M_2 = 150\text{GeV}$, $\mu = -400\text{GeV}$, and $\tan\beta = 2$.

as required to explain the HERA anomaly. However, scharm has also R conserving decay modes which are

$$\tilde{c}_L \rightarrow s\tilde{W}_i, s\tilde{Z}_i \quad (13)$$

We have seen that the corresponding branching fraction B for the decay modes eq. 12 is $B \leq 1/20$. Therefore, the direct decay channel eq. 12 will not give any viable SUSY signal. We have considered the cascade decays of the squarks and gluinos into the LSP which gives one electron through its decay as

$$\tilde{Z}_1 \xrightarrow{\lambda'_{121}} \bar{c}e^+d(\bar{s}\bar{\nu}d) + h.c. \quad (14)$$

As a consequence, the pair of LSP provides like sign dilepton(LSD) signature in this case due to its Majorana nature. We have considered the all possible strong production process¹⁰ like pair of gluino, pair of squark and associated production⁸ and their subsequent decays according to the masses of electroweak gauginos⁸. Considering only the λ'_{121} as leading RPV coupling, as suggested by the scharm scenario, we have estimated the LSD rates for a given value of charm squark mass(=210 GeV) applying some kinematic cuts

to suppress the SM background. We have seen that one can expect to see at least half of dozen LSD events with the present Tevatron luminosity of 110 pb^{-1} . Finally, we conclude with an expectation that the Tevatron group will analyse their data to probe the RPV SUSY model, and in particular to test the charm squark scenario for the anomalous HERA events.

The work reported here is done with collaboration of D.P.Roy⁸.

References

1. H1 Collaboration: C. Adloff et.al, *Z. Phys.* **C74**, 191 (1997).
2. ZEUS Collaboration: J. Breitweg et.al, *Z. Phys.* **C74**, 207 (1997).
3. G. Wolf, hep-ex/9704006; see however M. Drees, hep-ph/9703332
4. D. Choudhury and S. Raychaudhuri, hep-ph/9702392 and hep-ph/9703369.
5. H. Dreiner and P. Morawitz, hep-ph/9703279 (version 2).
6. G. Altarelli, J. Ellis, G.F. Giudice, S. Lola and M.L. Mangano, hep-ph/9703276.
7. S. Belayev and A.V. Gladyshev, hep-ph/9703251 and hep-ph/9704343; J. Kalinowski, R. Ruckl, H. Spiesberger and P.M. Zerwas, hep-ph/9703288; K.S. Babu, C. Kolda, J. March-Russell and F. Wilczek, hep-ph/9703299; V. Barger, K. Cheung, K. Hagiwara and D. Zeppenfeld, hep-ph/9703311.
8. M. Guchait and D.P. Roy, hep-ph/9707275 (to be published in Physical Review D).
9. H.P. Nilles, *Phys. Rep.* **110**, 1 (1984); H. Haber and G. Kane, *Phys. Rep.* **117**, 75 (1985); S. Weinberg, *Phys. Rev.* **D26**, 287 (1982); N. Sakai and T. Yanagida, *Nucl. Phys.* **B197**, 83 (1982); S. Dimopoulos, S. Raby and F. Wilczek, *Phys. Lett.* **B212**, 133 (1982).
10. G.L. Kane and J.P. Leveille, *Phys. Lett.* **B112**, 227 (1982); P.R. Harrison and C.H. Llewellyn-Smith, *Nucl. Phys.* **B213**, 223 (1983) [Err. *Nucl. Phys.* **B223**, 542 (1983)]; S. Dawson, E. Eichten and C. Quigg, *Phys. Rev.* **D31**, 1581 (1985); E. Reya and D. P. Roy, *Phys. Rev.* **D32**, 645 (1985).
11. CDF Collaboration: C. Grosso-Pilcher, Talk at the Vanderbilt Conference, May 1997.
12. DØ Collaboration: D. Norman, Talk at the Hadron Collider Physics Conference, Stonybrook, June 1997.
13. M. Guchait and D.P. Roy, *Phys. Rev.* **D54**, 3276 (1996).
14. H. Baer, C. Kao and X. Tata, *Phys. Rev.* **D51**, 2180 (1995).